

Ribbon structure in symmetric pre-monoidal categories

Liam Wagner¹, Jon Links² and Phil Isaac³

Centre for Mathematical Physics, The University of Queensland, Brisbane, 4072, Australia

Abstract. Let $U(g)$ denote the universal enveloping algebra of a Lie algebra g . We show the existence of a ribbon algebra structure in a particular deformation of $U(g)$ which leads to a symmetric pre-monoidal category of $U(g)$ -modules.

1. Introduction

Recently there has been some work [1, 2, 3, 4] on investigating the properties of categories in which the pentagon axiom which is central to the notion of monoidal categories [5] is forsaken. In particular it was shown in [6] that a certain deformation of the Hopf algebra structure of the universal enveloping algebras of Lie algebras naturally gives rise to a construction for symmetric pre-monoidal categories as defined in [2, 3, 4]. It is then reasonable to ask to what extent can such categories also be endowed with a ribbon structure, in analogy with the known examples of ribbon categories arising from representations of ribbon quasi-Hopf algebras [7]. Here we note that in a particular specialisation of the deformation given in [6] one can indeed preserve the ribbon algebra structure which provides an initial insight into making the notions of ribbon and pre-monoidal categories compatible.

2. Symmetric pre-monoidal categories

We shall begin by defining a pre-monoidal category, taken from [6].

Definition 1. A pre-monoidal category is a triple $(\mathcal{C}, \otimes, \mathcal{A})$ where $\mathcal{C}$ is the category of objects, $\otimes$ is a bifunctor $\otimes : \mathcal{C} \otimes \mathcal{C} \rightarrow \mathcal{C}$ and $\mathcal{A}$ is the natural associator isomorphism such that $\mathcal{A} : \otimes(\text{id} \times \otimes) \rightarrow \otimes(\otimes \times \text{id})$.

¹ ldw@maths.uq.edu.au

² jrl@maths.uq.edu.au

³ psi@maths.uq.edu.au

In practice we write $a_{U,V,W}$ as the action of the associator such that

$$a_{U,V,W} : U \otimes (V \otimes W) \rightarrow (U \otimes V) \otimes W.$$

We note that there are no conditions imposed on $\mathcal{A}$, but it is important that we should define the natural isomorphism $Q : \otimes(\otimes \times \otimes) \rightarrow \otimes(\otimes \times \otimes)$ via the following diagram:

$$\begin{array}{ccc}
(U \otimes V) \otimes (W \otimes Z) & \xleftarrow{q_{U,V,W,Z}} & (U \otimes V) \otimes (W \otimes Z) \\
\downarrow a_{(U \otimes V),W,Z} & & \uparrow a_{U,V,(W \otimes Z)} \\
((U \otimes V) \otimes W) \otimes Z & & U \otimes (V \otimes (W \otimes Z)) \\
\uparrow a_{U,V,W} \otimes \text{id} & & \downarrow \text{id} \otimes a_{V,W,Z} \\
(U \otimes (V \otimes W)) \otimes Z & \xleftarrow{a_{U,(V \otimes W),Z}} & U \otimes ((V \otimes W) \otimes Z)
\end{array}$$

This box diagram expresses Q in terms of the associator isomorphisms which are defined as:

$$Q((U \otimes V) \otimes (W \otimes Z)) = q_{U,V,W,Z}((U \otimes V) \otimes (W \otimes Z)). \quad (1)$$

These may be expressed as

$$q_{U,V,W,Z} = a_{(U \otimes V),W,Z}^{-1} (a_{U,V,W} \otimes \text{id}) a_{U,(V \otimes W),Z} (\text{id} \otimes a_{V,W,Z}) a_{U,V,(W \otimes Z)}^{-1}. \quad (2)$$

This condition is a generalisation of the pentagon condition used in describing monoidal categories, where $Q = \text{id} \otimes \text{id} \otimes \text{id} \otimes \text{id}$. We can examine the significance of Q and its use in distinguishing the coupling of the objects of the category through employing distinct brackets $[], \{\}$. This notation shows that

$$Q([U \otimes V] \otimes \{W \otimes Z\}) = (\{U \otimes V\} \otimes [W \otimes Z]). \quad (3)$$

We describe pre-monoidal categories as being unital if they have an identity object $1 \in \mathcal{C}$ and natural isomorphisms $\rho_U : U \otimes 1 \rightarrow U$ and $\lambda_U : 1 \otimes U \rightarrow U$. The next important class of unital pre-monoidal categories is when the tensor product is commutative up to isomorphism. This leads to the concept of a braided unital pre-monoidal category.

Definition 2. A unital pre-monoidal category $\mathcal{C}$ is said to be braided if it is equipped with a natural commutativity isomorphism $\sigma_{U,V} : U \otimes V \rightarrow V \otimes U$ for all objects $U, V \in \mathcal{C}$ such that the following diagrams commute [4]:

(i)

$$\begin{array}{ccccc}
& & U \otimes (V \otimes W) & \xrightarrow{\sigma_{U,(V \otimes W)}} & (V \otimes W) \otimes U \\
& \nearrow a_{U,V,W}^{-1} & & & \searrow a_{V,W,U}^{-1} \\
(U \otimes V) \otimes W & & & & V \otimes (W \otimes U) \\
& \searrow \sigma_{U,V} \otimes \text{id} & & & \nearrow \text{id} \otimes \sigma_{U,W} \\
& & (V \otimes U) \otimes W & \xrightarrow{a_{V,U,W}^{-1}} & V \otimes (U \otimes W)
\end{array}$$

(ii)

$$\begin{array}{ccccc}
& & (U \otimes V) \otimes W & \xrightarrow{\sigma_{(U \otimes V), W}} & W \otimes (U \otimes V) \\
& \nearrow a_{U, V, W} & & & \searrow a_{W, U, V} \\
U \otimes (V \otimes W) & & & & (W \otimes U) \otimes V \\
& \searrow \text{id} \otimes \sigma_{V, W} & & & \nearrow \sigma_{U, W} \otimes \text{id} \\
& & U \otimes (W \otimes V) & \xrightarrow{a_{U, W, V}} & (U \otimes W) \otimes V
\end{array}$$

(iii)

$$\begin{array}{ccc}
(U \otimes V) \otimes (W \otimes Z) & \xrightarrow{q_{U, V, W, Z}} & (U \otimes V) \otimes (W \otimes Z) \\
\downarrow \sigma_{(U \otimes V), (W \otimes Z)} & & \downarrow \sigma_{(U \otimes V), (W \otimes Z)} \\
(W \otimes Z) \otimes (U \otimes V) & \xleftarrow{q_{W, Z, U, V}} & (W \otimes Z) \otimes (U \otimes V)
\end{array}$$

Hereafter we will only be concerned with the case of *symmetric* categories in which the commutativity isomorphism satisfies the additional condition $\sigma_{U, V} \circ \sigma_{V, U} = \text{id}_{V \otimes U}$ for all objects $U, V \in \mathcal{C}$, and $\circ$ denotes composition of morphisms. In this case the diagrams (i) and (ii) are equivalent.

3. Twining

It is known that the finite-dimensional modules of quasi-triangular quasi-bialgebras give rise to braided monoidal categories, and that these categories are invariant under twisting [8, 9, 10]. Here we need to use a different method to build pre-monoidal categories. We follow the work of [6] where a twining operation was introduced to achieve this.

Define A as a quasi-triangular quasi-bialgebra by the octuple $(A, \Delta, \varepsilon, \Phi, \mathcal{R}, S, \alpha, \beta)$ where $\Delta : A \rightarrow A \otimes A$ is the co-product, Φ is an invertible element called the co-associator which is defined as $\Phi = I \otimes I \otimes I$, $\varepsilon : A \rightarrow \mathbb{C}$ is the co-unit, $\mathcal{R} \in A \otimes A$ is the universal R -matrix satisfying

$$\mathcal{R}\Delta(x) = \Delta^T(x)\mathcal{R} \quad \forall x \in A$$

where Δ^T is the opposite co-product. Further S is the antipode and α, β are canonical elements satisfying certain properties (see [8, 9, 10] for details). Let K be an element of the centre of A and let non-zero $\gamma \in \mathbb{C}$ be fixed but arbitrary. Following [6] we define

$$\begin{aligned}
\tilde{\mathcal{R}} &= \gamma^{K \otimes K} \mathcal{R} = \mathcal{R} \cdot \gamma^{K \otimes K} \\
\tilde{\Phi} &= \Phi \cdot \gamma^\kappa
\end{aligned} \tag{4}$$

such that $\kappa = K \otimes (I \otimes K + K \otimes I - \Delta(K))$. In general we write

$$\begin{aligned}
\tilde{\Phi} &= \sum_j X_j \otimes Y_j \otimes Z_j \\
\tilde{\Phi}^{-1} &= \sum_j \bar{X}_j \otimes \bar{Y}_j \otimes \bar{Z}_j \\
\tilde{\mathcal{R}} &= \sum_j a_j \otimes b_j.
\end{aligned} \tag{5}$$

Then the following hold:

$$\begin{aligned}
(\text{id} \otimes \Delta)\Delta(a) &= \tilde{\Phi}^{-1}(\Delta \otimes \text{id})\Delta(a)\tilde{\Phi} \quad \forall a \in A, \\
\tilde{\mathcal{R}}\Delta(a) &= \Delta^T(a)\tilde{\mathcal{R}}, \\
(\Delta \otimes \text{id})\tilde{\mathcal{R}} &= \tilde{\Phi}_{312}\tilde{\mathcal{R}}_{13}\tilde{\Phi}_{132}\tilde{\mathcal{R}}_{23}\tilde{\Phi}_{123}^{-1}, \\
(\text{id} \otimes \Delta)\tilde{\mathcal{R}} &= \tilde{\Phi}_{213}^{-1}\tilde{\mathcal{R}}_{13}\tilde{\Phi}_{213}^{-1}\tilde{\mathcal{R}}_{12}\tilde{\Phi}_{123}(\gamma^{2\kappa})_{123}^{-1}
\end{aligned} \tag{6}$$

and we go on to define

$$\xi = (\Delta \otimes \text{id} \otimes \text{id})\tilde{\Phi}^{-1}(\tilde{\Phi} \otimes I) \cdot (\text{id} \otimes \Delta \otimes \text{id})\tilde{\Phi} \cdot (I \otimes \tilde{\Phi}) \cdot (\text{id} \otimes \text{id} \otimes \Delta)\tilde{\Phi}^{-1}. \tag{7}$$

These relations show that the category $\mathbf{mod}_K(A)$ of A -modules with

$$a_{U,V,W} = (\pi_U \otimes \pi_V \otimes \pi_W)\tilde{\Phi}, \tag{8}$$

is a pre-monoidal category as $\tilde{\Phi}$ fails the pentagon condition. In particular the representation

$$q_{U,V,W,Z} = (\pi_U \otimes \pi_V \otimes \pi_Z)\xi \tag{9}$$

does not act as the identity.

We should note that $\tilde{\mathcal{R}}$ cannot be used in general as an ingredient in the construction of a braided, pre-monoidal category of A -modules, as the $\gamma^{2\kappa}$ defined above in (4) leads to a violation of the hexagon condition (i) of Definition 2. However, it was observed in [6] that in the case of the universal enveloping algebra $U(g)$ of a Lie algebra g , it is possible to choose $\gamma = -1$ and $K \in A$ taking integer eigenvalues on all irreducible finite-dimensional $U(g)$ -modules such that $\mathbf{mod}_K(A)$ does possess the structure of a symmetric pre-monoidal category. For a given central element $K \in U(g)$ such that $\varepsilon(K) = 0$ we call $(U(g), \Delta, \varepsilon, \tilde{\Phi}, \tilde{\mathcal{R}})$ with $\tilde{\Phi}$ and $\tilde{\mathcal{R}}$ given by (4) a *twining* of $U(g)$.

4. Ribbon algebra structure

In analogy with the definition of ribbon quasi-Hopf algebras [7], we can also investigate the ribbon structure for twined algebras. First recall that each quasi-triangular quasi-Hopf algebra A possesses a distinguished invertible element u satisfying

$$S^2(a) = uau^{-1} \quad \forall a \in A, \tag{10}$$

$$S(\alpha)u = \sum_j S(b_j)\alpha a_j. \tag{11}$$

We then have from [7]:

Definition 3. *Let A be a quasi-triangular quasi-Hopf algebra. We say that A is a ribbon quasi-Hopf algebra if there exists a central element $v \in A$ such that*

$$i. \ v^2 = uS(u)$$

$$ii. \ S(v) = v$$

iii. $\varepsilon(v) = 1$

iv. $\Delta(uv^{-1}) = \mathcal{F}^{-1}(S \otimes S)\mathcal{F}_{21}(uv^{-1} \otimes uv^{-1})$

where $\mathcal{F}$ is the Drinfeld twist [8] defined by the condition

$$\Delta(a) = \mathcal{F}^{-1}((S \otimes S)\Delta^T(S^{-1}(a)))\mathcal{F} \quad \forall a \in A.$$

The universal enveloping algebra $U(g)$ of a Lie algebra g acquires the structure of a quasi-bialgebra with mappings:

$$\begin{aligned} \varepsilon(I) &= 1, \quad \varepsilon(x) = 0, \quad \forall x \in g \\ S(I) &= I, \quad S(x) = -x, \quad \forall x \in g \\ \Delta(I) &= I \otimes I, \quad \Delta(x) = I \otimes x + x \otimes I, \quad \forall x \in g \end{aligned} \tag{12}$$

that are extended to all $U(g)$ such that ε and Δ are algebra homomorphisms and S is an anti-automorphism. It is easily checked that Δ is co-associative; i.e.

$$(\text{id} \otimes \Delta)\Delta(x) = (\Delta \otimes \text{id})\Delta(x) \quad \forall x \in U(g)$$

and co-commutative

$$\Delta(x) = \Delta^T(x) \quad \forall x \in U(g)$$

and that $S^2 = \text{id}$. This means that we can equip $U(g)$ with the structure of a quasi-triangular quasi-Hopf algebra by taking $\Phi = I \otimes I \otimes I$ for the co-associator of $U(g)$, $\mathcal{R} = I \otimes I$ as the universal R -matrix and $\alpha = \beta = I$. Note that in this instance the Drinfeld twist of $U(g)$ is trivial, and $U(g)$ trivially satisfies the conditions of a ribbon Hopf algebra with the choice $u = v = I$.

Under twining by a central element K satisfying $S(K) = -K$ we then have from (5)

$$\begin{aligned} \tilde{\mathcal{R}} &= (-1)^{K \otimes K} \\ \tilde{\Phi} &= (-1)^\kappa \end{aligned}$$

and moreover it can be shown that $\alpha = \beta = I$ (cf. [6]). It is easily verified that the choice

$$v = u = (-1)^{-K^2}$$

satisfies equations (10) and (11) as well as the conditions of Definition 3. Thus in this instance we can conclude that the twined algebra $U(g)$ can still be considered as a ribbon algebra.

5. Conclusion

We have shown that a particular case of the twining deformation described in [6] is compatible with the notion of a ribbon structure which can be endowed on quasi-triangular quasi-Hopf algebras. In future work we will explore the compatibility of ribbon categories and symmetric pre-monoidal categories on a general level and in particular investigate the consequences of this for defining traces and inner products on generic symmetric pre-monoidal categories (cf. [11]).

References

- [1] Yanofsky N S, Obstructions to coherence: natural noncoherent associativity 2000 *J. Pure Appl. Algebra* **147** 175
- [2] Joyce W P, Vertex Calculus in Premonoidal Categories: I. The Racah-Wigner Calculus, Recoupling and Statistics 2002 *Preprint*
- [3] Joyce W P, Quark State Confinement as a Consequence of the Extension of the Boson/Fermion Recoupling to $SU(3)$ Colour 2003 *J. Phys. A: Math. Gen.* **36** 12329
- [4] Joyce W P, Braided Premonoidal Mac Lane Coherence 2004 *J. Pure Appl. Algebra* **190** 155
- [5] Mac Lane S, 1998 *Categories for the Working Mathematician* 2nd. ed. (Springer-Verlag, New York)
- [6] Isaac P S, Joyce W P and Links J, An Algebraic Origin for Quark Confinement 2003 *Preprint* arXiv:hep-th/0307046
- [7] Altschuler D and Coste A, Quasi-quantum groups, knots, three-manifolds and topological field theory 1992 *Commun. Math. Phys.* **150** 83
- [8] Drinfeld V G, Quasi-Hopf algebras 1990 *Leningrad Math J.* **1** 1419
- [9] Chari V C and Pressley A, 1994 *A Guide to Quantum Groups* (Cambridge University Press)
- [10] Turaev V, 1994 *Quantum invariants of knots and 3-manifolds*, (W. de Gruyter, Berlin)
- [11] Kirillov A A, On Inner Product In Modular Tensor Categories, 1996 *J. Amer. Math. Soc.* **9** 155